Date: 1st February, 2024

ISSN: 2835-3196 Website: econferenceseries.com

PAIN IN THE BLADDER - SYMPTOMS, GENERAL INFORMATION

Xamidova Zilola Kamolova Tojigul Teachers of the Technical College of Public Health named after Abu Ali ibn Sino

Abstract:

The urinary bladder (bladder) is one of the main organs of the body's excretory system; we can say that it functions as a bag for storing fluid. Bladder pain can be caused by many different factors. For example, kidney disease, sexual dysfunction, inflammation in the buttocks. Such pain may indicate the presence of serious diseases not only of the bladder, but also of the organs located next to it.

About bladder pain

BLADDER PAIN

Bladder pain is a pathological symptom characterized by the appearance of various pain sensations in the lower abdomen. This is always accompanied by the quality and quantity of urine. All this allows us to call it a pain syndrome, since in this case it has its own characteristics, the details of which make it possible to determine the cause of the pain. In fact, pain is not an independent disease, but is a symptom that can be found in many diseases; it acts as a signal that you should be concerned. The adequacy of treatment depends on how correctly these data are interpreted.

The characteristics of pain in the bladder include the following: the nature of the pain, intensity, conditions of occurrence, what causes it, when it decreases, how long it lasts, exact localization and distribution, and other accompanying symptoms. In addition, it is necessary to take into account currently existing pathologies and past diseases, medical and other manipulations.

According to statistics, such pain is more often observed in women, and this is due to the structural features of their body. In women, the urinary tract is located in such a way that it is easier for them to become infected. Such diseases develop especially often in pregnant women, since in this case significant changes occur in the body's defense system and it becomes more difficult to fight the infection.

Cystitis is usually caused by various staphylococci and bacteria or becomes inflamed due to damage to the lining of the bladder. The feeling of pain can be a consequence of medical procedures, chemical or thermal burns. Acute or chronic cystitis can be





Proceedings of International Conference on Educational Discoveries and Humanities Hosted online from Plano, Texas, USA.

Date: 1st February, 2024

ISSN: 2835-3196 Website: econferenceseries.com

detected in pregnant women. If the disease is not treated, the inflammation begins to spread and affects the kidneys and urinary tract.

IN MEN

Such pain in men is a consequence of problems in the reproductive and reproductive systems. If there are stones or inflammation in the kidneys, additional symptoms may occur - diarrhea, increased body temperature and fever. In this case, you should consult a doctor. If the presence of a kidney stone is not detected, then the cause of pain in the bladder is urinary retention. To solve the problem, it is necessary to determine the factor that caused this delay.

Prostate adenoma often manifests itself in the form of pain in the lower abdomen. Usually men do not immediately realize the existence of this disease, since it occurs in a latent form. The disease can cause problems in the vascular system, disturbances in the circulatory system, and bother the patient under stress. The pain can be localized in the lower abdomen, in the groin area, in the testicles, and urination is always accompanied by pain. In this case, it is necessary to quickly make a diagnosis and prescribe the correct treatment methods.

Among the causes of pain in the bladder are the following diseases:

Acute and chronic cystitis is an inflammatory lesion of the walls of the bladder. Urethritis is an inflammation of the urinary tract.

Stones localized in the bladder.

Dyshormonal circulatory disorders of a menopausal nature in the wall of the bladder.

Tumors and polyps of the bladder wall.

Traumatic injury to the bladder and urethra.

Urolithiasis is accompanied by renal colic, when the stone moves from the kidney through the urinary tract to the bladder.

Prostatitis and benign prostatic hyperplasia (adenoma).

Inflammatory and tumor diseases of the uterus and appendages.

Diffuse pain due to injuries of the pubic symphysis, intestines and spine.

References:

- 1. A. Ahmedov, G. Ziyamutdinova: Anatomy, physiology and pathology.
- 2. Nerves and stress.
- 3. www.ziyonet.uz
- 4. https://mymedic.uz/

