

Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

MORPHOLOGICAL STRUCTURE OF INTESTINAL LAMBALISM, SYMPTOMS OF LAMBLIOSIS, DIAGNOSIS AND TREATMENT METHODS

DJumanova Nargiza Eshmamatovna,
Toshmamatov Bakhtiyar Norbekovich
Samarkand State Medical University

Norimov son of Azimjon Qahramon
2 Year Student

Annotation

Occurs in the form of vegetative as well as cysts in the cycle of life. The vegetative form is non-existent, 12-15km long and 8-10km wide. On the front side of the body is a disc that serves to attach to the intestinal epithelium. In the middle of the body are two base threads — axostils. Using axostils, the body of the parasite is divided into two wire metric nymphs, each with one nucleus. At the front of the axostils are several blepharoplasts, which provide energy to 3 pairs of chives. The fourth pair of chicks lies on the back of the axostils. The parasite does not have digestive organs, it is fed by an osmotic path. The method of reproduction is non-sexual, the path of division into vows is accomplished by mitosis. The cysts of the lambs are oval in shape and range in length from 10-14 μm . The two-story shell is clearly visible. Immature cysts have 2 nuclei and 4 nuclei when mature. The development cycle of lamblia is simple, and the period of life consists of the exchange of vegetative and cystic forms of the parasite. Vegetative forms live and multiply in a person's duodenum. In animals infected by experimental (experimental) pathways, lamblids are found in the upper parts of the small intestine. Some of them are wrapped in cysts, released into the external environment, and remain a source of human transmission of lambliosis. The source of lamblioz infection is the patient and the parasite carriers. Symptoms of the disease are similar to those of other intestinal diseases. Therefore, the main indicator of diagnosing lyambliosis should be the detection of parasites. But it's a much harder job. Twelve-fingered intestinal fluid contains vegetative forms of the parasite. The patient is diagnosed with parasitic cysts by iodine staining of the litter. With their rubbish, lamed cysts go out into the external environment and can survive for weeks. To a healthy person, the parasite is



Proceedings of International Educators Conference

Hosted online from Rome, Italy.


Date: 25th March, 2024

ISSN: 2835-396X


Website: econferenceseries.com

transmitted by cysts through contaminated food, drinking water, or dirty hands. After the cysts enter the digestive tract, the twelve fingers become a vegetative form in the intestine.

Keywords: Lyambliosis, parasitic cysts, diarrhea, epilepsy, lamblia intenstinalis, boss.



1. Intestinal lyambliosis in the intestine.



2. Microscopic morphology of intestinal lyambliosis.

Proceedings of International Educators Conference


Hosted online from Rome, Italy.

Date: 25th March, 2024


ISSN: 2835-396X

Website: econferenceseries.com

Relevance: Lamblia is ubiquitous and can now be found in 10 percent of the population. Lyambliosis is more common in children, mainly in children aged 1-4 years. Parasites can live in the gallbladder as well as in the duodenum. It can also be conditionally pathogenic, as the parasite is also found in healthy people, only when their number increases in the gut, intestinal activity changes. It can also be said to be unmarked, in the form of diarrhea, liver and bile duct disease. Absorption of substances as a result of impaired intestinal function decreases, adversely affecting the exchange of vitamins. This disease disrupts the functioning of the food system: the child is upset, the abdomen is heavy, the appetite is lost, and at the same time, the breakdown of nutrition causes a lot of disease, malnutrition due to loss of appetite can reduce the fight and cause seizures in the most severe complications.



3. intestinal placement and morphology of intestinal lyambliosis.



4. intestinal placement and morphology of intestinal lyambliosis.



Proceedings of International Educators Conference


Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

The purpose of the work: to gain an understanding of the forms (types) of Lyambliosis disease and to study the application of this disease to the sick in practice, having studied in theory the causes of common origin and more recent methods of treatment measures.



5. Microscope view of the cyst of intestinal lyambliosis.

Research Findings: The pathogenetic effects of lyamblia are as follows, and parasites disrupt the digestion and absorption of food in the gut. Sometimes diarrhea can also occur in the form of liver and biliary tract diseases. Loss of appetite, difficulty having breakfast in the morning, walking on an empty stomach all day, upper abdomen hurts, abdomen rests, collapses, nausea (yellow is a slightly slimy mixture), diarrhea o 'self-departure, thin stooling can sometimes cause seizures, sometimes the temperature can rise. A patient with this disease is treated for a long time. Symptomatic (unmarked) cyst toxicity is often observed. Laboratory diagnosis is made of these test methods. Through a method of iodine dyeing of feces, we identified their cystic forms in the litter and used parasite-lowering drugs. Of the drugs, we mainly used the drug Albindazole. We used Albindazole and a number of other similar drugs to lower the lyamblia. The basis for treating lambiosis is to follow a personal gegiena. In the treatment of lyamblia, the drug that kills him is taken for 5 days, but the liver should be thoroughly examined before giving this drug. Then



Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024


ISSN: 2835-396X

Website: econferenceseries.com

the dead lizards are lowered into the 5-day drink. If 20ku or 1oy grass driver (corn popup) is given, the lumbar will die.



6. Morphology of intestinal trichomonads.




7. Structure and parts of intestinal trichomoniasis.

Ways of washing

If there is no good destruction of feces, feces can contaminate water sources and food. Similarly, non-compliance with normal hygiene habits, such as not washing hands after using the bath, also means a common source of contamination. Not



washing fruits, vegetables, greens well can also cause infection and is also contagious when eating with unwashed hands. Mosquitoes can serve as mechanical transmission factors as well as crowded and very close connections.



8. Intestinal lamblia infection and development cycle.

Trophozoites or cysts should be monitored in fecal samples, duodenal juice, or biopsies to detect the parasite.

Methods of disease prevention:

It focuses mainly on a person's personal hygiene.

- Thorough washing of raw fruits, vegetables and other ingredients;
- boiling water and drinking;
- Cooked food and drinking water in a closed container save;



Proceedings of International Educators Conference


Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

- shaving; washing hands before and after meals;
- Includes tasks such as timely detection and treatment of parasitic carriers, especially those working in food enterprises and kindergartens



9. Microscope view of the intestinal lily.

Treatment

The drugs selected for intestinal lyambliya are quinacrine hydrochloride or nitroimidazoles. Among nitroimidazoles:

- Metronidazole (50 mg / kg per day, divided into 3 doses from 7 to 10 days).
- Tinidazole (60 mg / kg per day for 1 mg to 3 days).

Furazolidone is often used in pediatric patients because it is present in liquid suspension, but the treatment rate is lower.

Due to the risk of teratogenicity, none of the above-mentioned drugs can be used in pregnant women.

The only drug recommended for pregnant women is paromomycin, which is safer because it is not absorbed, even if it is less effective

Conclusion:

In short, the disease occurs in almost every country, and in patients with this disease is characterized by impaired intestinal digestion and absorption of food. The aim of



Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

this study is to study this common disease in more depth, to study the causes of origin, diagnostics and treatment in more depth, to create new methods of treatment and to put it into practice. A person with lyamblia is required to undergo laboratory examination in consultation with a pediatrician, parasitologist, until the lyamblia is completely removed from the body. It should be borne in mind that the main reason for the origin of this disease is the non-compliance with personal hygiene and public hygiene in the lifestyle, and in conclusion, most people follow the rules of hygiene to prevent the disease, everyone is healthy at least 2 times in 1 year. we ask the doctor to go and get a checkup so that we can not only pre-determine other diseases like this, but now prevent treatment at the initial time.

References:

1. Parasitology: A learning hand for students of pedagogical universities./S.Dadayev; Ministry of Higher and Secondary Special Education of the Republic of Uzbekistan. - T.: «Platinum Publishers», 2006. - 208 p.
2. Djumanova N. E., Qaxramon o'g'li A. N. SALTING OF SUCKERS AND THE DISEASES THEY CAUSE LIVERWORM //World Bulletin of Public Health. – 2023. – T. 21. – C. 82-85.
3. Djumanova N. E., Murodullayeva B. K. et al. PROPERTY DISTROFIY DISEASE, SIMPTOMS, DIAGNOSTICS, METHODS OF IMPROVEMENT VALUATION // FANLARO INNOVATIONS AND SCIENTIFIC RESEARCH IN UZBEKISTAN. – 2023. – 2. – №. 17. – S. 220-224.
4. Djumanova N. E., Murodullayeva B. K. et al. PROPERTY DISTROFIY DISEASE, SIMPTOMS, DIAGNOSTICS, METHODS OF IMPROVEMENT VALUATION // FANLARO INNOVATIONS AND SCIENTIFIC RESEARCH IN UZBEKISTAN. – 2023. – 2. – №. 17. – S. 220-224.
5. Dzhumanova N. E., Nazarova F. S. PROBABLE NEGATIVE IMPACT OF GENETICALLY MODIFIED PRODUCTS ON HUMAN HEALTH //Thematics Journal of Botany. – 2022. – T. 6. – №. 1.
6. Dzhumanova N. E., Nazarova F. S. PROBABLE NEGATIVE IMPACT OF GENETICALLY MODIFIED PRODUCTS ON HUMAN HEALTH //Thematics Journal of Botany. – 2022. – T. 6. – №. 1.
7. Nishonboyev K.N, Hamidov J.H. Daxlik. - Tashkent: Biology State Scientific Publishing House, 2005. - 210 pages.



Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

8.Nazarova F.Sh., Khudoiberdieva G., Dzhumanova N.E. Biochemical comparative analysis of the ecological composition of phytonematodes.

9.Nazarova F. Sh., Dzhumanova N. E., Donierova P. H. DIAGNOSIS OF HEREDITY AND SYMPTOMS OF IMPAIRED CARBOHYDRATE METABOLISM AND GALACTOSEMIA. – 2023.

10.Nazarova F. Sh., Dzhumanova N. E. The use of bentonite from the Azkamara deposit for balancing mineral nutrition // Academic research in the field of pedagogical sciences. – 2021. – Issue 2. – No. 9. – С. 672-679.

11.Nazarova F.C., Dzhumanova N.E. Hair and wool cover as indicators of environmental pollution by technogenic and geochemical sources // Thematic Journal of Microbiology. – 2022. – Issue 6. – No. 1.

12.Nazarova F.S., Dzhumanova N.E., Tashmamatov B.N., Sh. O. Korzhavov. The phytonematode ecological group. Problems of biology and medicine. - 2020. No. 6. Volume 124. - С.

13.Nazarova F.S., Dzhumanova N.E. The biological role of trace elements and their content in epidermal formations. Economy and society.1-2 (92).2022. pp. 94-103.

14.Назарова Ф. Ш., Джуманова Н. Э., Дониёрова П. Х. ДИАГНОСТИКА НАСЛЕДСТВЕННОСТИ И СИМП-ТОВ НАРУШЕ-НИЯ образовательного процесса и ГАЛАКТОЗЕМИИ //ИННОВАЦИИ В ТЕХНОЛОГИЧЕСКОМ И ЕСТЕСТВЕННОНАУЧНОМ ОБРАЗОВАНИИ. – 2023. – Т. 2. – №. 7. – С. 446-454.

15.Nazarova F. S., Eshmamatovna Djumanova N., Danielova P. H. DETERMINATION OF CARBONSUV ALMACHINE AND IRSY OF GALACTOZEMIA DISEASE, DIAGNOSTICS OF SIMPTOMS //Innovations in Technology and Science Education. – 2023. – Т. 2. – №. 7. – S. 438-445.

16.Nazarova F.Sh., Khudoyberdiyeva G., Djumanova N.E. Biochemical sravnitelny analysis ekologicheskogo sostava phytonematode.

17.Nazapova F. Sh., Djumanova N. E. Icpolzovanie bentonita azkamapckogo mestopojdeniya dlya balancipovki minepalnogo pitaniya / / Akademicheskie icledovaniya v oblacti pedagogicheckix nauk. – 2021. - VIP. 2. – № 9. – С. 672-679.

18.Nazapova F.C., Djumanova N.E. Volocno-shepctniy pokpov kak indicatopi zagpyazneniya okpuyayushey cpedi technogennimi I geochemicheckimi ichtochnikami // tematichescki jupnal mikpobiologii. – 2022. - VIP. 6. – №1.

19.Nazapova F.C., Djumanova N.E., Tashmamatov B.N., Sh. O. Kopjavov.



Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

Ecologicheckaya guppipovka phytonematode. Poblemi biologii I medicini. - 2020. № 6. Tom 124. - С. 258-261.

20. Najapova F.Sh., Dzhumanova N.E. Biological Paul of trace elements and their codeposition in epidermal formations. Economy and society.1-2 (92).2022. стр. 94-103.

21. Nazarova F. S., Dzhumanova N. E. HAIR AND WOOL AS INDICATORS OF ENVIRONMENTAL POLLUTION BY MAN-MADE AND GEOCHEMICAL SOURCES //Thematics Journal of Microbiology. – 2022. – Т.6.– №. 1.

22. Nazarova F. S. et al. BIOLOGICAL ROLE OF MICROELEMENTS AND THEIR CONTENT IN EPIDERMAL FORMATIONS //European Journal of Molecular and Clinical Medicine. – 2021. – Т. 8. – №. 2. – S. 16751687.

23. Nazarova F. S., Dzhumanova N. E. HAIR AND WOOL AS INDICATORS OF ENVIRONMENTAL POLLUTION BY MAN-MADE AND GEOCHEMICAL SOURCES //Thematics Journal of Microbiology. – 2022. – Т. 6. – №. 1.

24. Тошмаматов Б.Н. и соавт. Макрокопическая схема илеоцекального локута у полицейских //Международная научно-практическая конференция «Мировая наука. – РОСТ, 2017. – С. 5. – № 5. – С. 58-59.

25. Тошмаматов Б.Н. и соавт. Макроскопическая структура илеоцекального локута у кроликов //Международная научно-практическая конференция «Мировая наука. – РОСТ, 2017. – С. 5. – № 5. – С. 58-59.

26. Toshmamatov B. N. et al. MACROSCOPIC COMPARISON OR OCULAR REJECTION IN RATS AND RABBITS IN STATE ONTOGENESIS //International scientific and practical conference World science. – GROWTH, 2018. – vol. 3. – No. 5. – pp. 53-54.

27. Toshmamatov B.N. i soavt. Макроскопическая структура илеоцекального отвёртиа кролика i кроликов v ростнатальном ontogeneze //Международная научно-практическая конференция «Мировая наука. – РОСТ, 2018. – № 3. – № 5. – С. 53-54.

28. Toshmamatov B.N. Морфологические изменения желудка при полипрагмазии Домашняя страница журнала Science LLC Поступила в редакцию 04.04.2021 Поступила в редакцию 20.04.2021ю

29. Toshmamatov B.N., Tashaev Sh.J., Khudoiberdiev D.K. Consequences of polypragmasia with preventive drugs on the coprology of the tenka-stomach, problems of biology and medicine 2022, No.1 (134) С.157-164.



Proceedings of International Educators Conference

Hosted online from Rome, Italy.

Date: 25th March, 2024

ISSN: 2835-396X

Website: econferenceseries.com

30. Tokhmetov B.N., Tataev V. O., Khudoiberdiev V.L., Morphological assessment of the climatic lining of the stomach in case of polyppagmasia with anti-inflammatory drugs // Russian Scientific Journal, Volume 20, January 2022, pages 32-38.

31. Tomatov B.N., Tanaev Yu. A change in the morphological structure of the enema membrane in polyppagmasia. (2021). American International Medical Journal of Scientific and Pharmaceutical Research, 3 (08), 51-57.

32. Toshmamatov B.N., Tshaev Sh.Dj., Xudoybepdiev D.K. Izmenenie morfologicheskix pokazateley clizicnoy obolochki jeludka i osnovaniya clizicnoy jeludka belqx krcs ppi polippagmazii. Problemy biologii i meditsiny 6. 2022, №3 (136) С.190 -195.

33. Tshaev Sh.J., B.N. Toshmamatov Gistologicheskie izmeneniya jeludka ppi lechenii nosteroidnymi ppreparatami Problemy biologii i meditsiny 2 (34) 2021/ С.260-264.

34. Toshmamatov B. N. i dr. Makroskopicheskoye stroeniye ileotsekal'noy zaslonki u krolikov. //International Scientific and Practical Conference World science. – ROST, 2017. – Т. 5. – №. 5. – S. 5859.

35. Toshmamatov B. N. i dr. Makroskopicheskoye stroeniye ileotsekal'nogo otverstiya u krolikov i krys v postnatal'nom ontogeneze //International Scientific and Practical Conference World science. – ROST, 2018. – Т. 3. – №. 5. – S. 53-54.

36. Тошмаматов Б. Н. и др. МАКРОСКОПИЧЕСКОЕ СТРОЕНИЕ ИЛЕОЦЕКАЛЬНОЙ ЗАСЛОНКИ У КРОЛИКОВ //International Scientific and Practical Conference World science. – ROST, 2017. – Т. 5. – №. 5. – С. 58-59.

