

EKSTREMAL MASALALARNI YECHISHDA TENGSIZLIKLAR

USULIDAN FOYDALANISH

Raximova Muxtaram Rashidovna

Qo‘qon DPI Akademik litseyi
matematika fani bosh o‘qituvchisi

Annotatsiya

Ushbu maqolada Ekstremal masalalarni yechishda tengsizliklar usulidan foydalanish hususida alohida to‘xtalib o‘tilgan. Shuningdek, maqolada tengsizliklar haqida ham ayrim misollar keltirilgan.

Kalit so‘zlar. Tengsizlik, ekstremal, usul, jihat, yechish, hosila, mohiyat, ko‘nikma

ИСПОЛЬЗОВАНИЕ МЕТОДА НЕРАВЕНСТВ ПРИ РЕШЕНИИ

ЭКСТРЕМАЛЬНЫХ ЗАДАЧ.

Рахимова Мухтарам Рашидовна

Коканский ГПИ Академический Лицей
старший преподаватель математики

Абстрактный

В данной статье основное внимание уделяется использованию метода неравенств при решении экстремальных задач. В статье также приведены некоторые примеры неравенств.

Ключевые слова. Неравенство, крайность, метод, аспект, решение, производная, суть, умение.

USING THE METHOD OF INEQUALITIES IN SOLVING EXTREME PROBLEMS

Rakhimova Mukhtaram Rashidovna

Kokan SPI Academic Lyceum
head teacher of mathematics



Abstract

This article focuses on the use of the method of inequalities in solving extreme problems. The article also provides some examples of inequalities.

Keywords. Inequality, extreme, method, aspect, solution, derivative, essence, skill

Kirish

Tengsizliklar usuli va uning afzalliklari haqida. Ekstremal masalalarni yechishning turli usullari mavjud. Bunga avvalo, hosilalardan foydalanib yechish usulini (hosila usulini) misol sifatida keltirish mumkin. Qolaversa, geometrik usul xamda tengsizliklar usuli ham shular jumlasidandir. Biz bu maqolada ba'zi muhim klassik tengsizliklarning ekstremal masalalar yechishga tatbig'i (tengsizliklar usuli) bilan shug'ullanamiz. Ekstremal masalalarni yechishda tengsizliklar usulining mohiyati quyidagicha bayon etlishi mumkin.

Masalaning qo'yilishi. Ushbu maqolada quyidagi masalani tadqiq etamiz. Ixtiyoriy musbat x_1, x_2, \dots, x_k sonlar berilgan bo'lsin. Ular yordamida ushbu

$A_k = \frac{x_1 + x_2 + \dots + x_k}{k}$, $\Gamma_k = \sqrt[k]{x_1 x_2 \dots x_k}$ kattaliklarni tuzamiz. Bunda A_k – berilgan

sonlarning o'rta arifmetigi, Γ_k esa ularning k – tartibli o'rta geometrigi deyiladi.

Mazkur kattaliklar orasida $\Gamma_k \leq A_k$ (1)

munosabat mavjud bulib, unda tenglik belgisiga $x_1 = x_2 = \dots = x_k$ bo'lganda va faqat shu holdagina erishiladi.

Masalaning yechilishi Ushbu $x_1 + x_2 + \dots + x_k$ kattalik o'zgarmas bo'lsin, uni a bilan belgilaylik. U holda (1) tengsizlikdan ko'rinadiki, Γ_k ning eng katta qiymati

faqat $A_k = \frac{a}{k}$ dan iborat bo'lishi mumkin. Bunga esa, $x_1 = x_2 = \dots = x_k$

bo'lgandagina erishiladi. Demak, k ta musbat son yig'indisi o'zgarmas bo'lsa, shu sonlar ko'paytmasidan chiqarilgan k — darajali ildiz o'zining eng katta qiymatiga

ular o'zaro teng bo'lgandagina erishadi va bu eng katta qiymat $A_k = \frac{a}{k}$ ning o'zi

bo'ladi. Shunday qilib, $\max \Gamma_k = \frac{a}{k}$.



2) Endi $x_1 x_2 \dots x_k$ kattalik o'zgarmas son deylik, uni b bilan belgilaymiz. U holda (1) tengsizlikdan ravshanki, A_k ning eng kichik qiymati $\sqrt[k]{b}$ bo'lishi mumkin holos. Bunga esa $x_1 = x_2 = \dots = x_k$ bo'lgandagina erishiladi. Demak k ta musbat son ko'paytmasi o'zgarmas bo'lsa, shu sonlarning o'rta arifmetigi o'zining eng kichik qiymatiga ular o'zaro teng bo'lgandagina erishadi va bu eng kichik qiymat $\Gamma_k = \sqrt[k]{b}$ ning o'zi bo'ladi. Shunday qilib, $\min A_k = \sqrt[k]{b}$.

Tengsizliklar usuli ma'lum sinf ekstremal masalalarni yechishda hosila usuliga qaraganda ba'zi afzalliklarga ega:

- 1) ekstremal qiymati topilishi kerak bo'lgan funksiyaning hosilasi hisoblanmaydi;
- 2) Topilishi kerak bo'lgan ekstremal qiymat (ekstremal hajm, yuza, uzunlik, burchak, vaqt va hokazolar) darhol topiladi;
- 3) Shu ekstremal qiymatga erishiladigan nuqta (hosila usulidagi kritik nuqtalar ichida funksiyaga ekstremal qiymat beradigani) ham darhol topiladi;
- 4) Tengsizliklar usuli ekstremal qiymati topilishi lozim bo'lgan funksiya ko'p argumentli bo'lganda ham qo'llanilishi mumkin.

2. Tengsizliklardan geometriya kursi va ekstremal masalalarni yechishda foydalanish. Endi (1) tengsizlikni avval planimetriyadagi, so'ngra stereometriyadagi ba'zi ekstremal masalalarni yechishga tatbiq etamiz. Bu maqolada planimetriyaga oid quyidagi masala ko'ramiz.

Masalaning amaliy tadbirlari. Asosi c va balandligi h bo'lgan uchburchakka bir tomoni asosda, ikki uchi uchburchakning yon tomonlarida bo'lgan eng katta yuzali to'g'ri to'rtburchak chizilsin.

Yechish: Chizmada $\triangle ABC$ va $\square A_1, B_1, B_2, A$ lar chizilgan. Unda $AB = c$, $CD = h$ deylik. Ushbu $A_1 B_1 = A_2 B_2 = x$, $A_1 A_2 = ED = B_1 B_2 = y$ belgilashlarni kiritamiz.

$\triangle ABC$ va $\square A_2, B_2, C, D$ larning o'hshashligidan $\frac{c}{h} = \frac{x}{h-y}$, bundan $x = \frac{c}{h}(h-y)$

$A_1 B_1 B_2 A_2$ to'g'ri to'rtburchakning yuzi $S(y) = xy = \frac{c}{h} y(h-y)$, ammo



$y(h-y) \leq \left(\frac{h}{2}\right)^2$ va bu tengsizlikda tenglik « \Rightarrow » belgisi $y = h-y$ bo'lganda



erishiladi. U holda $y = \frac{h}{2}$, $x = \frac{c}{2}$. Demak, $S_{\max} = \frac{c}{h} \cdot \frac{h^2}{4} = \frac{1}{2} \cdot c \cdot \frac{h}{2} = \frac{1}{2} S_{\Delta ABC}$.

Shunday qilib, to'g'ri to'rtburchakning A_2B_2 tomoni uchburchak balandligining o'rtasidan uchburchak asosiga parallel qilib o'tkazilsa, ya'ni A_2B_2 chiziq ΔABC ning o'rta chizig'i bo'lsa, to'g'ri to'rtburchakning yuzi eng katta bo'ladi.

2-masala. Radiusi R bo'lgan aylana ichiga chizilgan teng yonli uchburchaklar ichida yuzi eng kattasi topilsin.



Yechish. Chizmada, $AC + CB$, $OA = OB$, $CD = x$. Ravshanki,

$$DO = x - R, AB = 2AD = 2\sqrt{R^2 - DO^2} = 2\sqrt{R^2 - (x - R)^2} = 2\sqrt{2Rx - x^2}.$$

Endi $S_{\Delta ABC}$ yuzini topamiz. $S_{\Delta ABC} = S(x) = \frac{1}{2} \cdot x \cdot \sqrt{2Rx - x^2} = \sqrt{x^3} \sqrt{2R - x}$.

Unda $0 < x < 2R$. Qiymatini topish uchun tengliklar usulini to'g'ridan to'g'ri qo'llab bo'lmaydi. $S(x)$ ni kvadratga ko'taramiz: $S^2(x) = x^3(2R - x)$

va bu funktsiyani quyidagicha yozib olamiz: $S^2(x) = \frac{1}{3} \cdot x \cdot x \cdot x \cdot (6R - 3x)$

Agar $x_1 = x, x_2 = x, x_3 = x, x_4 = 6R - 3x$ deb belgilasak

$$x_1 + x_2 + x_3 + x_4 = 6R = \text{const}$$

va $x = 6R - 3x$, $x = \frac{3}{2}R$ kelib chiqadi. Demak, tengsizliklar usulini qo'llash

mumkin. $\Gamma_4 \leq A_4$ tengsizlikka ko'ra

$$S^2(x) = \frac{1}{3} \cdot x \cdot x \cdot x \cdot (6R - 3x) \leq \frac{1}{3} \left(\frac{x + x + x + 6R - 3x}{4} \right)^4 = \frac{1}{3} \left(\frac{6R}{4} \right)^4 = \frac{1}{3} \left(\frac{3}{2}R \right)^4$$

Bundan $S(x) \leq \frac{1}{\sqrt{3}} \left(\frac{3}{2}R \right)^2 = \frac{9}{4\sqrt{3}} R^2$, ya'ni $S_{\max} = S\left(\frac{3}{2}R\right) = \frac{9}{4\sqrt{3}} R^2$

kelib chiqadi. Endi ΔABC ning tomonlarini topib qo'yaylik:

$$AB = 2\sqrt{2R \cdot \frac{3}{2}R - \left(\frac{3}{2}R\right)^2} = 2\sqrt{3R^2 - \frac{9}{4}R^2} = \sqrt{3R},$$

$$AC - CB - \sqrt{AD^2 + DC^2} = \sqrt{\left(\frac{AB}{2}\right)^2 + x^2} = \sqrt{\frac{3}{4}R^2 + \frac{9}{4}R^2} = \sqrt{3R}$$

Shunday qilib, $AB = BC = CA = \sqrt{3R}$. Bu natija $\triangle ABC$ ning teng tomonli ekanligi anglatadi. Shunday qilib, radiusi R bo'lgan aylana ichiga chizilgan teng yonli uchburchaklar ichida yuzi eng kattasi teng tomonli uchburchak bo'lar ekan.

Xulosa

Ushbu maqolada ko'rilgan masalalar matematikaning eng muhim masalalaridan biri hisoblanadi. Maqoladan mustaqil o'rganish va izlanishlar olib borish maqsadida akademik litsey, kasb-hunar kollejlari va oliy ta'lim muassasasi talabalari manba sifatida foydalanishlari mumkin.

Foydalanilgan adabiyotlar

1. F.B.Badalov, F.Shodmonov "Matematik modellar va muhandislik masalalarini sonli yechish usullari". "Fan" nashriyoti. Toshkent 2000 yil.
2. Хикматов. "Экстремал масалалар". Тошкент, 1989.
3. G'.Nasriddinov, X.Shokirova. SH.Shoziyotov. "Stereometriya kursida ekstremal masalalar".

